

A NOTE ON p -ADIC INVARIANT INTEGRAL IN THE RINGS OF p -ADIC INTEGERS

TAEKYUN KIM

*Jangjeon Research Institute for Mathematical Sciences and Physics,
252-5 Hapcheon-Dong Hapcheon-Gun Kyungnam, 678-802, S. Korea
e-mail: tkim@kongju.ac.kr/ tkim64@hanmail.net*

ABSTRACT. In [2], I constructed the p -adic q -integral $I_q(f)$ on $\mathbb{Z}_p$. In this paper, we consider the properties of $\lim_{q \rightarrow -1} I_q(f) = I_{-1}(f)$. Finally we give the some applications of $I_{-1}(f)$ and integral equations for $I_{-1}(f)$. These properties are useful and worthwhile to study Euler numbers and polynomials.

1. INTRODUCTION

Let p be a fixed odd prime. Throughout $\mathbb{Z}_p$, $\mathbb{Q}_p$ and $\mathbb{C}_p$ will respectively denote the rings of p -adic integers, the fields of p -adic numbers and the completion of of the algebraic closure of $\mathbb{Q}_p$. For d a fixed positive integer with $(p, d) = 1$, let

$$X = X_d = \varprojlim_N \mathbb{Z}/dp^N, \quad X_1 = \mathbb{Z}_p,$$

$$X^* = \bigcup_{\substack{0 < a < dp \\ (a, p) = 1}} a + dp\mathbb{Z}_p,$$

$$a + dp^N\mathbb{Z}_p = \{x \in X \mid x \equiv a \pmod{dp^N}\},$$

where $a \in \mathbb{Z}$ lies in $0 \leq a < dp^N$, (cf. [1], [2]).

1991 *Mathematics Subject Classification.* 11S80.

Key words and phrases. p -adic invariant integral, q -Volkenborn integral.

The p -adic absolute value in $\mathbb{C}_p$ is normalized so that $|p|_p = \frac{1}{p}$. Let q be variously considered as an indeterminate a complex number $q \in \mathbb{C}$, or a p -adic number $q \in \mathbb{C}_p$. If $q \in \mathbb{C}_p$, we assume $|q - 1|_p < 1$ for $|x|_p \leq 1$. Throughout this paper, we use the following notation :

$$[x]_q = [x : q] = \frac{1 - q^x}{1 - q} = 1 + q + q^2 + \cdots + q^{x-1}.$$

We say that f is uniformly differentiable function at a point $a \in \mathbb{Z}_p^-$ and denote this property by $f \in UD(\mathbb{Z}_p^-)$ if the difference quotients

$$F_f(x, y) = \frac{f(x) - f(y)}{x - y},$$

have a limit $l = f'(a)$ as $(x, y) \rightarrow (a, a)$, cf. [1, 2]. For $f \in UD(\mathbb{Z}_p)$, let us start with the expression

$$\frac{1}{[p^N]_q} \sum_{0 \leq j < p^N} q^j f(j) = \sum_{0 \leq j < p^N} f(j) \mu_q(j + p^N \mathbb{Z}_p), \text{ cf. [1, 2],}$$

representing q -analogue of Riemann sums for f .

The integral of f on $\mathbb{Z}_p$ will be defined as limit ($n \rightarrow \infty$) of these sums, when it exists. An invariant p -adic q -integral of a function $f \in UD(\mathbb{Z}_p)$ on $\mathbb{Z}_p$ is defined by

$$I_q(f) = \int_X f(x) d\mu_q(x) = \int_{\mathbb{Z}_p} f(x) d\mu_q(x) = \lim_{N \rightarrow \infty} \frac{1}{[p^N]_q} \sum_{0 \leq j < p^N} f(j) q^j, \text{ see [2].}$$

Note that if $f_n \rightarrow f$ in $UD(\mathbb{Z}_p)$; then

$$\int_{\mathbb{Z}_p} f_n(x) d\mu_q(x) \rightarrow \int_{\mathbb{Z}_p} f(x) d\mu_q(x).$$

The classical Euler polynomials are defined by

$$F(t, x) = \frac{2}{e^t + 1} e^{xt} = \sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!}.$$

Note that $E_n = E_n(0)$ are called n -th Euler numbers. In [3], an analogue of Bernoulli number is defined by $\frac{t}{we^t - 1} = \sum_{n=0}^{\infty} B_n(w) \frac{t^n}{n!}$, where w is the element of locally constant functions' space [3]. For $f \in UD(\mathbb{Z}_p)$, we define a p -adic invariant integral on $\mathbb{Z}_p$ as follows:

$$I_{-1}(f) = \lim_{q \rightarrow -1} I_q(f) = \int_{\mathbb{Z}_p} f(x) d\mu_{-1}(x).$$

The purpose of this paper is to give the integral equations related to $I_{-1}(f)$ and to investigate some properties for $I_{-1}(f)$. From these integral equations, we derive the interesting formulae related to Euler numbers.

§2. p -ADIC INVARIANT INTEGRALS ON $\mathbb{Z}_p$

For $f \in UD(\mathbb{Z}_p)$, the p -adic q -integral was defined by

$$(1) \quad I_q(f) = \lim_{N \rightarrow \infty} \frac{1}{[p^N]_q} \sum_{0 \leq x < p^N} q^x f(x) = \lim_{N \rightarrow \infty} \sum_{0 \leq x < p^N} f(x) \mu_q(x + p^N \mathbb{Z}_p) = \int_{\mathbb{Z}_p} f(x) d\mu_q(x),$$

representing p -adic q -analogue of Riemann integral for f (see [2]). In the meaning of fermionic, we now define $I_{-1}(f)$ -integral as

$$(2) \quad I_{-1}(f) = \lim_{q \rightarrow -1} I_q(f) = \lim_{q \rightarrow -1} \int_{\mathbb{Z}_p} f(x) d\mu_{-q}(x) = \int_{\mathbb{Z}_p} f(x) d\mu_{-1}(x).$$

From (2), we derive the below Eq.(3):

$$(3) \quad I_{-1}(f) = \int_{\mathbb{Z}_p} f(x) d\mu_{-1}(x) = \lim_{N \rightarrow \infty} \sum_{x=0}^{p^N-1} (-1)^x f(x).$$

Let $f_1(x) = f(x+1)$. Then we note that

$$I_{-1}(f_1) = - \lim_{N \rightarrow \infty} \sum_{x=0}^{p^N-1} f(x) (-1)^x + 2f(0) = -I_{-1}(f) + 2f(0).$$

Therefore we obtain the below theorem:

Theorem 1. *For $f \in UD(\mathbb{Z}_p)$, we have*

$$(4) \quad I_{-1}(f_1) + I_{-1}(f) = 2f(0),$$

where $f_1(x)$ is translation with $f_1(x) = f(x+1)$.

From Theorem 1, we can derive the below theorem:

Theorem 2. *For $f \in UD(\mathbb{Z}_p)$, $n \in \mathbb{N}$, we have*

$$I_{-1}(f_n) + (-1)^{n-1} I_{-1}(f) = 2 \sum_{x=0}^{n-1} f(x) (-1)^{n-1-x},$$

where $f_n(x) = f(x+n)$.

By using Theorem 1 and Theorem 2, we can consider the new extension of Euler numbers and polynomials. If we take $f(x) = \lambda^x e^{xt}$, ($\lambda \in \mathbb{Z}_p$), then we have

$$(5) \quad \frac{2}{\lambda e^t + 1} = \int_{\mathbb{Z}_p} \lambda^x e^{xt} d\mu_{-1}(x).$$

Define the new extension of Euler numbers as follows:

$$(6) \quad \frac{2}{\lambda e^t + 1} = \sum_{n=0}^{\infty} E_n(\lambda) \frac{t^n}{n!}.$$

By (5) and (6), we obtain the below theorem:

Theorem 3. (Witt's formula) For $n \in \mathbb{N}$, we have

$$\int_{\mathbb{Z}_p} \lambda^x x^n d\mu_{-1}(x) = E_n(\lambda), \lambda \in \mathbb{Z}_p,$$

where $E_n(\lambda)$ are called analog of Euler numbers.

By using I_{-1} -integral, we define the new extension of classical Euler polynomials as follows:

$$(7) \quad \int_{\mathbb{Z}_p} \lambda^y (x+y)^n d\mu_{-1}(y) = E_n(\lambda : x).$$

From (7), we can derive the below:

$$(8) \quad E_n(\lambda : x) = \sum_{k=0}^n \binom{n}{k} E_n(\lambda) x^{n-k} = f^n \sum_{a=0}^{f-1} (-1)^a \lambda^a E_n(\lambda^f : \frac{x+a}{f}),$$

where f is odd positive integer.

By (8), we easily see that

$$(9) \quad \frac{2e^{xt}}{\lambda e^t + 1} = \int_X e^{(x+y)t} d\mu_{-1}(y) = \frac{2 \sum_{a=0}^{f-1} (-1)^a \lambda^a e^{(x+a)t}}{\lambda^f e^{ft} + 1}.$$

Remark 1. In [3], it was known that

$$(9-1) \quad \lim_{q \rightarrow 1} \int_X w^y e^{(x+y)t} d\mu_q(t) = \frac{te^{xt}}{we^t - 1} = \sum_{m=0}^{\infty} B_m(w) \frac{t^m}{m!},$$

where $B_m(w)$ are called an analogue of Bernoulli numbers. In the viewpoint of (9-1), we gave the new extension $(E_n(\lambda))$ of classical Euler number.

Remark 2. From using multivariate p -adic invariant integral for $I_{-1}(f)$, we can easily derive the Euler polynomials of higher order as follows:

$$e^{xt} \int_{\mathbb{Z}_p} \dots \int_{\mathbb{Z}_p} \lambda^{y_1 + \dots + y_r} e^{(y_1 + \dots + y_r)t} d\mu_{-1}(y_1) \dots d\mu_{-1}(y_r) = \left(\frac{2}{\lambda e^t + 1} \right)^r e^{xt}.$$

From this, we can define the extension of classical Euler polynomials of order r as follows:

$$\left(\frac{2}{\lambda e^t + 1} \right)^r e^{xt} = \sum_{n=0}^{\infty} E_n^{(r)}(\lambda : x).$$

Let χ be the Dirichlet's character with conductor $f(= \text{odd}) \in \mathbb{N}$ and let us take $f(x) = \chi(x)e^{tx}$. From Theorem 2, we derive the below formula:

$$(10) \quad \int_X e^{tx} \chi(x) d\mu_{-1}(x) = \frac{2 \sum_{a=0}^{f-1} e^{ta} (-1)^a \chi(a)}{e^{ft} + 1}.$$

The generalized Euler numbers attached to χ were defined by

$$(11) \quad \frac{2 \sum_{a=0}^{f-1} e^{ta} (-1)^a \chi(a)}{e^{ft} + 1} = \sum_{n=0}^{\infty} E_{n,\chi} \frac{t^n}{n!}, \text{ cf. [4].}$$

From (10) and (11), we derive the below Witt's formula for the generalized Euler numbers attached to χ as follows:

Theorem 4. Let f be an odd positive integer and let χ be the Dirichlet's character with conductor f . Then we have

$$\int_X x^n \chi(x) d\mu_{-1}(x) = E_{n,\chi}.$$

REFERENCES

- [1] T. Kim, *q*-Volkenborn integration, Russ. J. Math. Phys. **9** (2002), 288-299.
- [2] T. Kim, *On a q-analogue of the p-adic log gamma functions and related integrals*, J. Number Theory **76** (1999), 320-329.
- [3] T.Kim, *An analogue of Bernoulli numbers abd their congruences*, Rep. Fac. Sci. Engrg. Saga Univ. Math. **22** (1994), 21-26.
- [4] T.Kim, *p-adic l-functions and sums of powers*, arXiv:math.NT/0605703 **1** (2006).